

Supplemental Table S1. Comparison of the patients with or without postoperative ileus

	Postoperative ileus (+) (n = 7)	Postoperative ileus (-) (n = 41)	P-value
Age (yr)	62.86±14.51	63.10±10.53	0.958
Male sex	5 (71.4)	25 (61.0)	0.501
BMI (kg/m ²)	20.16±2.34	22.89±3.12	0.033
ASA grade 3	2 (28.6)	7 (17.1)	0.662
Emergency operation	1 (14.3)	5 (12.5)	0.643
Previous admission history	1.71±1.50	4.43±7.41	0.344
Time to bowel frequency (POD, day)	11.57±5.00	8.50±9.10	0.391
Fever	3 (42.9)	13 (31.7)	0.447
Leukocytosis	5 (71.4)	12 (29.3)	0.049
Admission duration (POD, day)	15.71±20.08	7.62±3.88 ^{a)} 8.04±3.92 ^{b)}	0.328 0.078
Treatment for CDI (yes)	3 (42.9)	31 (75.6)	0.080
Death	1 (14.3)	2 (4.9)	0.391

Values are presented as mean ± standard deviation or number (%).

BMI, body mass index; ASA, American Society of Anesthesiologists; POD, postoperative days; CDI, *Clostridium difficile* infection.

^{a)}Resection group+colectomy group. ^{b)}Colectomy group only.